SEX-AGE CONSTITUTION AND DIAGNOSIS OF THE 317 NEW PATIENTS DURING 1914.

	1 —			5 —			10 —			15 —			25 —			35 —			45 —			55 —			65 +			All Ages		
	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes TOTAL
—	9	3	—	9	14	—	5	8	—	13	16	—	14	15	—	9	10	—	10	7	—	3	—	—	—	—	—	72	73	145
—	2	1	—	3	3	—	1	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	6	6	12
—	7	15	—	12	8	—	12	8	—	3	6	—	4	13	—	—	9	—	2	8	—	—	3	—	—	—	—	45	72	117
—	1	4	—	2	9	—	4	2	—	2	5	—	2	6	—	—	4	—	1	—	—	—	1	—	—	—	—	12	31	43
—	19	23	—	26	34	—	22	19	—	18	27	—	20	35	—	9	23	—	13	15	—	3	4	—	—	—	—	135	182	317

PERCENTAGE IN EACH AGE-GROUP FOUND TO HAVE PULMONARY TUBERCULOSIS.

	1 —			5 —			10 —			15 —			25 —			35 —			45 —			55 —			65 +			All Ages		
	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes TOTAL
—	19	23	—	26	34	—	22	19	—	18	27	—	20	35	—	9	23	—	13	15	—	3	4	—	—	—	—	135	182	317
—	9	3	—	9	14	—	5	8	—	13	16	—	14	15	—	9	10	—	10	7	—	3	—	—	—	—	—	72	73	145
—	47·36	13·04	—	34·61	41·17	—	22·72	42·1	—	72·2	59·25	—	70·0	42·85	—	100·0	43·47	—	76·92	46·6	—	100·0	—	—	—	—	—	53·3	40·1	45·7
—	2	1	—	3	3	—	1	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	6	6	12
—	10·52	4·34	—	11·53	8·82	—	4·54	5·26	—	—	—	—	—	2·85	—	—	—	—	—	—	—	—	—	—	—	—	—	4·4	3·29	3·78